



SOI FAX HOTLINES

(Southern Oscillation Index)

this page = 1902 935 301

this page is updated each Wednesday, usually by 5pm



SOI MESSAGE - 11 November 1998

SOI (a) AVERAGES / PHASE		
August 98	*	+9.8
September 98	*	+12.1
October 98	*	+11.2

Last 30 days		+ 15.3
Last 90 days (b)		+ 11.9

SOI trend (b) during September-October was Phase 2 (i.e. positive ▲).		
NOTE : (a) SOI values calculated using mean barometric pressures from 1880 to 1992 - subject to revision by Bureau of Meteorology. * Preliminary value ** Revised value (b) See AUSTRALIAN RAINMAN for effects of SOI on rainfall at your location.		

grain is about 15% moisture, followed by grain drying, is likely to pay dividends. Also in summer-cropping areas, where there is good soil moisture, consider planting some crop into winter crop residues after harvesting.

Colder-than-normal, equatorial sea-surface temperatures are now present from around the International Dateline to the eastern Pacific Ocean. In addition, temperatures are considerably warmer than normal in waters off northern Australia.

Above-average pasture growth is likely over much of Queensland during the November-January period. The favourable outlook may provide opportunities for burning native pasture, and sowing improved pastures.

The average SOI over the last 30 days has increased to +15.3 . The probabilities of exceeding median rainfall for the November to January period are 60% to 80% in most of Queensland and New South Wales.

THE BOTTOM LINE

REVIEW OF CLIMATIC FORECASTS AND INFORMATION

Most climatic indicators suggest continued high rainfall probabilities.

Use of the AUSTRALIAN RAINMAN package indicates that the probabilities of exceeding median rainfall during the November-January period are generally 60-80% in the eastern States of Australia. However, parts of north-western and central-western Queensland, south-western NSW, central and western Victoria, and Tasmania have probabilities of about 50%.

Some USA agencies are predicting the development of a La Niña pattern this year. Such a development would probably result in reduced rainfall for our trade competitors in south-western USA, Argentina and central Asia.

In the current situation, there is an above-average risk of rain interfering with winter crop harvesting, and causing losses. Commencing harvest when mature

A preliminary assessment indicates that the next passage of the 30- to 50-day Oscillation is due about the first week in December.

To obtain more detailed information for your location, we recommend combined use of the AUSTRALIAN RAINMAN package and the Bureau of Meteorology's Seasonal Climate Outlook. Also a lot of additional information is available on our SOI Fax Hotlines, our Internet World Wide Web service called 'The Long Paddock', and on BoM's fax and Internet information services.

NEXT UPDATE of the SOI MESSAGE:
18th November 1998

Climate Impacts and Grazing Systems - Department of Natural Resources

Compiled by Col Paull and Dr Roger Stone, QDPI.

If you would like any further information, please contact Col Paull on (07) 389 69587, or one of the climate extension officers at the DPI in Charters Towers, Emerald, Gympie, Kingaroy, Longreach, Mackay, Mareeba, Roma and Toowoomba.

Some information courtesy Bureau of Meteorology, CSIRO and National Oceanographic and Atmospheric Administration, USA