



SOI FAX HOTLINES

(Southern Oscillation Index)

this page = 1902 935 301

this page is updated each Wednesday, usually by 5pm



SOI MESSAGE - 26 May 1999

| SOI (a) AVERAGES / PHASE | | |
|---|---|-------|
| February 99 | * | +7.1 |
| March 99 | * | +7.6 |
| April 99 | * | +16.8 |
| ----- | | |
| Last 30 days | | + 0.1 |
| Last 90 days (b) | | + 9.0 |
| ----- | | |
| SOI trend (b) during March-April was Phase 4 (i.e. rising ↗). | | |
| NOTE : (a) SOI values calculated using mean barometric pressures from 1880 to 1992 - subject to revision by Bureau of Meteorology. * Preliminary value ** Revised value (b) See AUSTRALIAN RAINMAN for effects of SOI on rainfall at your location. | | |

Rainfall prospects over the winter cropping areas of central and southern Queensland are mostly average. The SOI Phase at the end of February indicated the likelihood of frosts (-2°C in the screen) later in the winter than normal over winter cropping areas of central Queensland, and around normal in southern Queensland and northern NSW. However, the frost risk is generally very low as lack of rain has prevented widespread early plantings. Burning wheat stubble before planting this year's crop may be necessary to control yellow spot.

The likelihood of above-average pasture growth during the May-July period is low for much of Queensland, as pasture growth has already reached its potential in many regions. However, recent rainfall has enhanced prospects for pasture growth in parts of central-western and southern Queensland.

The average SOI over the previous 30 days has fallen to +0.1. While a weakened La Niña pattern continues, the probabilities of exceeding median rainfall during the total period of May to July are 60-80% in the south-eastern quarter of Queensland and the north-eastern corner of NSW, and about normal values over most of Queensland.

THE BOTTOM LINE

REVIEW OF CLIMATIC FORECASTS AND INFORMATION

SOI climate forecasting systems indicate that rainfall probability values in most parts of the eastern States of Australia during the May to July period vary little from the long-term 'normal' values for that time of year. Exceptions are the south-eastern quarter of Queensland and adjacent border districts in NSW, and coastal districts in northern NSW and eastern Victoria where probabilities are 60-80%. However, in parts of far southern NSW, central and western Victoria, and far northern Tasmania the probabilities are 30-40%.

A 'weakened' La Niña pattern continues in the Pacific Ocean. In general terms, such a pattern means reduced rainfall for our trade competitors in south-western USA, Argentina and central Asia.

Colder-than-normal, equatorial sea-surface temperatures are now present from around the International Dateline to the far eastern Pacific Ocean. In addition, temperatures are warmer than normal in waters off most of the eastern coastline of Australia.

The next passage of the 30- to 50-day Oscillation is due about the middle of June.

To obtain more detailed information for your location, we recommend combined use of the AUSTRALIAN RAINMAN package and the Bureau of Meteorology's Seasonal Climate Outlook. Also a lot of additional information is available on our SOI Fax Hotlines, our Internet World Wide Web service called 'The Long Paddock', and on BoM's fax and Internet information services.

NEXT UPDATE of the SOI MESSAGE:
2nd June 1999

Climate Impacts and Grazing Systems - Department of Natural Resources

Compiled by Col Paull and Dr Roger Stone, QDPI.

If you would like any further information, please contact Col Paull on (07) 389 69587, or one of the climate extension officers at the DPI in Charters Towers, Emerald, Gympie, Kingaroy, Longreach, Mackay, Mareeba, Roma and Toowoomba.

Some information courtesy Bureau of Meteorology, CSIRO and National Oceanographic and Atmospheric Administration, USA